

In erratum

The authors of the article Effect of Two Contrasting Interventions on Upper Limb Chronic Pain and Disability: A Randomized Controlled Trial (Emil Sundstrup, MSc et al. Pain Physician 2014; 17:145-154) have requested the following text changes to their article:

Fig. 2. Change in chronic pain intensity (VAS scale 0 – 10) from baseline (0-wks) to follow-up (10-wks) with resistance training (full lines) and ergonomic training (dashed lines). Values are means (SE). *, **, *** Denotes greater reductions in pain with resistance training compared to ergonomic training (Post hoc test: $P < 0.01$, $P < 0.001$, $P < 0.0001$, respectively).

P value in the Results: "($P < 0.001$, $P < 0.05$, $P < 0.0001$, respectively)" should be changed to "($P < 0.001$, $P = 0.05$, $P < 0.0001$, respectively)"

P value in the Results: "($P < 0.001$, $P < 0.05$, $P < 0.0001$, respectively)" should be changed to "($P < 0.001$, $P = 0.05$, $P < 0.0001$, respectively)"

Figure text for Fig.2: "($P < 0.05$, $P < 0.01$, $P < 0.001$, respectively)" should be changed to "(Post-hoc test: $P < 0.01$, $P < 0.001$, $P < 0.0001$, respectively)"

Figure text for Fig.3: "($P < 0.05$)" should be changed to "(Post-hoc test: $P < 0.05$)"

Abstract:

In the Results: "($P < 0.001$, $P < 0.05$, $P < 0.0001$, respectively)" should be changed to "($P < 0.001$, $P = 0.05$, $P < 0.0001$, respectively)"

And in all places "ergonomics" should be changed to "ergonomic"

Figure 3 and 4 were reversed in the manuscript Effects of Dexmedetomidine Alone and in Combination with Tramadol or Amitriptyline in a Neuropathic Pain Model (Pain Physician/March/April 2014; 17:187-195).




Fig. 3. Time course effect of saline on paw withdrawal threshold in rats with CCI. Significant differences evoked by electronic von Frey test every 15 minutes up to 120 minutes ($n = 6-8$) between saline-treated group compared with Pre CCI value; *, $P < 0.05$.




Fig. 4. Time course effect of saline on the pain threshold in rats with CCI. Significant differences evoked by Randal and Sellito test every 15 minutes up to 120 minutes ($n = 6$) between saline-treated group compared with Pre CCI value; *, $P < 0.05$.